ISSN 1998 – 409X

REVIEW ARTICLE

Phytohormones from Soybean: A Novel Approach to Modulating Secondary Sexual Characteristics in Early Adolescence

Ashish Majumdar[1], Jitendra Patel[2], Surendra Saraf[1], Chandrashekhar Sahu[3]

[1]Department of Pharmacy, Columbia College of Pharmacy, Raipur, Chhattisgarh, India, [2]Department of Pharmacy, Datta Meghe College of Pharmacy, Adani Institute of Higher Education and Research, Deemed to be University, Sawangi (M), Wardha, Maharashtra, India, [3]Department of Pharmacy, RITEE College of Pharmacy, Raipur, Chhattisgarh, India

Abstract

Soybean-derived phytohormones, particularly isoflavones, such as genistein and daidzein, have garnered significant attention for their potential impact on human health, especially in modulating hormonal pathways during adolescence. This review explores the biochemical properties of soy phytohormones and their role in the modulation of secondary sexual characteristics, such as breast development and pubic hair growth, during early adolescence. By examining pre-clinical and clinical studies, we evaluate the influence of soy isoflavones on pubertal development and hormonal regulation. Despite some concerns regarding the estrogenic effects of phytohormones, available evidence suggests that moderate soy consumption may offer health benefits, such as improved cardiovascular health and reduced risks of hormonal imbalances. However, the variability in individual responses and the limitations of present studies call for further research to establish conclusive safety guidelines. The findings indicate the potential of soy-based dietary interventions for personalized nutrition strategies while underscoring the need for long-term studies to assess their efficacy and safety in adolescent populations.

Key words: Adolescence, daidzein, dietary interventions, estrogenic effects, genistein, health benefits, hormonal regulation, isoflavones, phytohormones, pubertal development, secondary sexual characteristics, soybean

INTRODUCTION

Secondary sexual characteristics are key markers of human development, particularly during adolescence, when profound physiological and psychological transformations occur. These changes are primarily regulated by hormones, such as testosterone and estrogen, which are produced endogenously. Recent research has highlighted the potential influence of external dietary factors, including phytohormones, on the timing and progression of these changes. Soybean-derived phytohormones, due to their structural and functional similarity to human sex hormones, are of significant interest in understanding their modulatory effects on hormonal pathways during adolescence.[1] Nanotechnology is an emerging trend nowadays for better absorption and efficacy. Phytohormones are also being incorporated in nanoformulation.[2] The isolation techniques also help the researcher determine the phytoconstituents in the extracts.[3] Preliminary phytochemical investigation gives a general idea to proceed with further investigation in the same plant.[4]

Overview of phytohormones in soybeans

Soybeans are a rich source of bioactive compounds known as isoflavones, including genistein and daidzein, which are classified as phytoestrogens due to their ability to bind to estrogen receptors alpha and beta (ERα and ERβ). Depending on the hormonal environment, concentration, and receptor subtype, these isoflavones exhibit agonistic and antagonistic effects. Genistein, in particular, demonstrates a higher affinity for ERβ, which has been implicated in regulatory and protective roles in various tissues.[5]

The biochemical properties of soybean phytohormones allow them to mimic or interfere with endogenous hormone

Address for correspondence:
Jitendra Patel, Datta Meghe College of Pharmacy, Datta Meghe Institute of Higher Education, deemed to be University, Wardha, India.
E-mail: jitendra.pharmacy@dmiher.edu.in

Received: 21-07-2026
Revised: 28-08-2026
Accepted: 09-09-2026

© 2026 Author(s). This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0)

signaling pathways. For instance, genistein and daidzein can influence the activity of aromatase enzymes, which convert androgens into estrogens, thereby indirectly modulating androgen levels. In addition, these compounds undergo gut microbial metabolism to form bioactive metabolites, such as equol, which may further enhance or alter their physiological effects.[6]

Emerging studies suggest that dietary phytohormones can influence the onset and progression of puberty, with potential implications for early or delayed development of secondary sexual characteristics. While their effects are not uniform and depend on dosage, individual microbiota, and genetic predispositions, phytohormones are increasingly recognized as significant modulators of hormonal activity during adolescence.[7]

Importance of secondary sexual characteristics in adolescence

Secondary sexual characteristics, including breast development in females, deep voice in males, and pubic hair growth in both genders, signify the biological maturation associated with adolescence. These changes are driven by the activation of the hypothalamic-pituitary-gonadal (HPG) axis, which orchestrates the production of sex steroids. Such characteristics are physiological markers and have profound social, psychological, and health implications.[8]

The timing of pubertal development is critical; deviations from the norm can lead to significant health risks. Early onset of puberty, known as precocious puberty, has been linked to increased risks of metabolic disorders, reproductive health issues, and psychosocial challenges. Conversely, delayed puberty can affect growth patterns, bone density, and overall development. External influences, including dietary components, such as soybean phytohormones, are thought to play a role in modifying these timelines, especially in populations with high soy consumption.[9]

Given their potential to interact with estrogen and androgen signaling pathways, soybean phytohormones may contribute to pubertal onset and progression variations. Understanding these effects is essential to evaluate the broader implications of soy-based diets during adolescence and to establish appropriate dietary guidelines.[10]

Scope and objectives of the review

This review comprehensively examines the role of soybean-derived phytohormones in influencing secondary sexual characteristics during adolescence. It focuses on these phytohormones' biochemical properties and mechanisms, particularly their interaction with human hormonal pathways. In addition, the review evaluates evidence from both pre-clinical and clinical studies regarding the effects of soy-derived phytohormones on pubertal development and the emergence of secondary sexual characteristics.

The discussion extends to the potential health implications, exploring the benefits and risks of soy consumption during this critical developmental phase while emphasizing safety and efficacy considerations. To provide a forward-looking perspective, the review also highlights existing research gaps and proposes future directions better to understand the role of phytohormones in human development.

By synthesizing diverse findings, the review offers a nuanced understanding of how soybean phytohormones might influence adolescent health and development. Moreover, it aims to inform dietary guidelines and public health policies that effectively incorporate these bioactive compounds into adolescent nutrition strategies.

BIOCHEMISTRY OF SOYBEAN PHYTOHORMONES

Soybean is a rich source of bioactive compounds that mimic human hormones, making it a focal point of research in endocrinology. These compounds predominantly isoflavones, such as genistein and daidzein, exhibit structural similarities to endogenous estrogens, enabling them to interact with human hormonal pathways. This section delves into the biochemical properties of these phytohormones, their mechanisms of action, and their interactions with androgen and ER.[11]

Key hormonal compounds in soybean (isoflavones, genistein, daidzein)

Isoflavones are the primary phytohormones in soybeans, with genistein and daidzein being the most studied. These compounds are classified as phytoestrogens due to their ability to bind to ERs. Genistein, a 4',5,7-trihydroxyisoflavone, has a strong affinity for ER beta (ERβ), whereas daidzein, structurally simpler, also interacts with ERs but exhibits slightly lower binding affinity.

Genistein and daidzein are metabolized by the gut microbiota into bioactive metabolites, such as equol and O-desmethylangolensin. These metabolites significantly enhance the biological activity of isoflavones, particularly in individuals who are "equol producers" (Clarke and Bailey, 2017). This metabolic variability partly explains the diverse physiological effects of soy-based diets in different populations.[12]

Mechanisms of action on endocrine pathways

Soybean phytohormones act through several mechanisms, primarily modulating the endocrine system by interacting with hormonal receptors and enzymes. Genistein and daidzein

mimic the structure of endogenous estrogens, allowing them to bind to ERα and ERβ competitively. Their binding affinity and resultant biological activity are context-dependent, functioning as agonists or antagonists based on hormonal concentrations and receptor subtype expression.

These isoflavones also influence the enzymatic pathways involved in hormone biosynthesis. For example, genistein inhibits aromatase, the enzyme responsible for converting androgens into estrogens, thereby altering the balance between these hormones. This modulation can affect puberty onset and the development of secondary sexual characteristics interacts with tyrosine kinases, further influencing cellular signaling pathways that regulate growth and development.[13]

Interaction with androgen and ERs

Phytohormones from soybeans display a dual affinity for estrogen and androgen receptors (ARs), enabling them to exert complex regulatory effects. Their selective interaction with ERβ over ERα suggests a protective role in tissues, such as the brain, bones, and cardiovascular system. By modulating ERβ activity, these compounds help balance the effects of endogenous estrogens, which can influence the timing and intensity of secondary sexual characteristic development during adolescence.[14]

The interaction with ARs is less well-characterized but significant. Studies suggest that isoflavones may reduce the activity of ARs by downregulating androgen signaling pathways. This effect can delay or suppress androgen-driven changes, such as increased body hair and voice deepening, particularly in males consuming high-soy diets.[15]

The biochemistry of soybean phytohormones reveals their potential as modulators of endocrine activity, with significant implications for adolescent development. By binding to estrogen and ARs and influencing hormonal pathways, compounds, such as genistein and daidzein demonstrate a capacity to impact secondary sexual characteristics. However, the extent of these effects varies based on individual metabolism, dietary patterns, and genetic predisposition. Further research is needed to clarify these interactions and establish guidelines for soy consumption during critical developmental periods.[16]

ROLE OF PHYTOHORMONES IN PUBERTAL DEVELOPMENT

Soybean-derived phytohormones, particularly isoflavones, such as genistein and daidzein, have garnered significant interest for their ability to influence human hormonal pathways. These bioactive compounds can modulate pubertal development by interacting with key endocrine receptors, leading to physiological changes in secondary sexual characteristics. This section explores their influence on early puberty markers, compares natural and soy-derived hormones, and evaluates gender-specific effects.[17]

Influence on early secondary sexual characteristics

The onset of secondary sexual characteristics, such as breast development in females and facial hair growth in males, is tightly regulated by hormonal signaling. Phytohormones from soybeans act as selective ER modulators (SERMs), binding to ERα and ERβ and exerting estrogen-like or anti-estrogenic effects depending on the hormonal milieu.[18]

Research suggests that isoflavones can mimic low-level estrogen activity, potentially accelerating breast tissue growth in pre-pubescent girls. However, their anti-estrogenic properties may also delay pubertal progression under certain conditions. A longitudinal study observed that soy-based infant formula, rich in phytoestrogens, correlated with subtle variations in pubertal timing, but the clinical significance remains debated.[19]

In males, phytohormones may attenuate androgenic activity by inhibiting enzymes, such as 5-alpha-reductase, which converts testosterone to dihydrotestosterone, a potent androgen responsible for secondary characteristics, such as voice deepening and body hair growth. This modulatory effect underscores their potential to influence gender-specific puberty trajectories.[20]

Comparative analysis: Natural versus soy-derived hormones

Natural endogenous hormones' physiological impact differs significantly from soy-derived phytohormones. Endogenous estrogens and androgens have high receptor-binding affinities and tightly regulated feedback mechanisms, ensuring precise control over pubertal events. In contrast, phytohormones exhibit weaker affinities for ERs and ARs, often acting as partial agonists or antagonists.[21]

Natural hormones are synthesized and metabolized within the endocrine system, while soy phytohormones enter the body through dietary intake. This introduces variability in bioavailability, which depends on individual factors, such as gut microbiota composition. For instance, the ability to produce equol, a daidzein metabolite, enhances phytoestrogen activity in some individuals, leading to stronger hormonal effects.[22]

Moreover, while endogenous hormones are finely balanced, phytohormones may exert biphasic effects, acting as estrogenic at low concentrations and anti-estrogenic at higher levels. This duality complicates the interpretation of their influence on pubertal development, necessitating more nuanced studies.[23]

Gender-specific effects of soy phytohormones

The effects of soy phytohormones are notably gender-specific due to differences in hormone receptor expression and baseline hormonal profiles. In females, phytoestrogens may modestly mimic estrogen activity, potentially promoting early breast development but showing limited impact on other estrogen-regulated processes. In males, however, these compounds may inhibit androgen signaling, leading to reduced expression of masculine traits, such as slower progression of muscle growth or voice changes.

Studies have also highlighted variability in response to phytohormones based on genetic and environmental factors. For instance, males consuming high-soy diets exhibited slight reductions in testosterone levels, though the effects were not clinically significant. In females, soy intake correlated with less severe premenstrual symptoms, suggesting a potential therapeutic role in hormonal modulation.

Soy-derived phytohormones play a complex role in pubertal development, influencing secondary sexual characteristics through interactions with endocrine pathways. Their effects vary based on hormonal context, gender, and individual metabolism. While promising as natural modulators of puberty, further research is essential to clarify their safety, efficacy, and long-term implications for adolescent health.[24]

EVIDENCE FROM PRE-CLINICAL AND CLINICAL STUDIES

Phytohormones derived from soybeans, primarily isoflavones, such as genistein and daidzein, have been extensively studied for their influence on hormonal modulation. This section delves into the findings from animal studies, human research, and the challenges in interpreting experimental outcomes to understand the role of soy-derived phytohormones in secondary sexual development and adolescent health.[25]

Animal studies on hormonal modulation by soy

Animal studies provide foundational insights into how phytohormones interact with endocrine systems. These studies have primarily focused on the estrogenic and anti-estrogenic effects of isoflavones, given their structural similarity to 17β-estradiol. In rodent models, soy isoflavones have been shown to modulate the HPG axis, impacting hormone secretion and receptor activation.

Neonatal exposure to genistein resulted in altered ovarian function and disrupted estrous cycles in female mice. These effects were attributed to the compound's ability to bind ERα and ERβ, which regulate reproductive physiology. Similarly, studies on male rodents revealed that dietary isoflavones could suppress testosterone levels by inhibiting key enzymes, such as 5-alpha reductase, thereby reducing androgen activity.

While these findings highlight the hormonal impact of soy phytohormones, their relevance to human physiology is limited by species-specific differences in metabolism and receptor sensitivity. For instance, rodents metabolize phytoestrogens differently, often leading to higher systemic bioavailability than humans.[26]

Human studies and observational data in adolescents

Human studies on soy-derived phytohormones have yielded mixed results, particularly in adolescents, where hormonal balance is critical for pubertal development. Observational studies and randomized controlled trials (RCTs) have explored their effects on secondary sexual characteristics and hormonal markers.

One notable study assessed the impact of soy-based infant formula on pubertal timing in a cohort of adolescents. The study found minor variations in breast development in females and testicular volume in males; however, these changes were within clinically normal ranges. Another RCT evaluated the effect of soy protein supplementation in adolescent boys, concluding that it did not significantly alter testosterone levels or secondary sexual characteristics.

Despite these findings, some observational studies suggest potential benefits of phytohormones in mitigating hormonal imbalances. Isoflavones have been linked to reduced severity of premenstrual symptoms and improved hormonal homeostasis in adolescent females. However, variability in individual responses, influenced by genetic factors, such as the ability to produce equol – a metabolite of daidzein – complicates generalizations.[27]

Limitations and variability in experimental outcomes

Several limitations hinder the interpretation of both pre-clinical and clinical studies on soy phytohormones. Key challenges include:

- Variability in dosage and bioavailability: Animal studies often use higher isoflavone doses than typically consumed in human diets, limiting their applicability.
- Differences in metabolism: Human metabolism of phytohormones depends on gut microbiota composition, which varies significantly across populations.
- Short study durations: Most human studies are limited to short durations, making it difficult to assess long-term effects on puberty and development.
- Confounding factors: Factors, such as diet, genetics, and environmental exposures can influence study outcomes, adding complexity to data interpretation.

These limitations underscore the need for more rigorous, long-term studies with standardized methodologies to clarify the role of soy phytohormones in hormonal modulation and adolescent health.

Evidence from pre-clinical and clinical studies provides valuable insights into the hormonal effects of soy-derived phytohormones. While animal models highlight potential mechanisms, human studies reveal nuanced effects influenced by individual variability and study design. Addressing present limitations is essential for developing clearer dietary recommendations and understanding the long-term implications of soy consumption in adolescence.[28]

HEALTH IMPLICATIONS OF SOY PHYTOHORMONES IN ADOLESCENCE

Soy-derived phytohormones, primarily isoflavones, such as genistein and daidzein, have garnered significant attention for their potential effects on adolescent health. These bioactive compounds, known for their estrogen-like properties, interact with the endocrine system, influencing hormonal regulation and developmental processes. This section explores the benefits, risks, and dietary implications of soy phytohormones during adolescence.[29]

Benefits in hormonal regulation and development

One of the most significant advantages of soy phytohormones is their ability to modulate hormonal activity, particularly during adolescence, a critical period of hormonal fluctuation and physical development. Isoflavones act as SERMs, binding to ERα and ERβ and exerting either estrogenic or anti-estrogenic effects depending on the hormonal milieu, as provided in Table 1.

Studies suggest that soy phytohormones can help regulate menstrual cycles in adolescent females by promoting hormonal balance. Isoflavones reduce estrogen dominance, mitigating symptoms, such as irregular menstruation and premenstrual syndrome. In addition, phytohormones have been associated with improved bone density, a crucial factor for adolescents as they undergo rapid skeletal growth.

For adolescent males, soy phytohormones may have a stabilizing effect on testosterone levels, preventing the overproduction of androgens. The role of isoflavones in supporting hormonal homeostasis without significantly disrupting androgenic activity. Such modulation is particularly beneficial in preventing early or delayed onset of secondary sexual characteristics.[30]

Risks and concerns: Hormonal imbalance or overconsumption

While the benefits of soy phytohormones are noteworthy, concerns regarding their potential risks cannot be overlooked. Excessive consumption of soy isoflavones has been linked to hormonal imbalances, particularly in sensitive populations. Isoflavones' estrogenic activity raises concerns about their potential to disrupt endocrine functions.

In females, excessive soy intake may lead to delayed pubertal onset or hypoestrogenism, characterized by insufficient estrogen production. In males, high isoflavone consumption has been associated with reduced testosterone levels, potentially affecting sperm production and fertility. However, these effects are generally observed at consumption levels significantly higher than those found in typical diets.

Another concern is the variability in individual responses to phytohormones, influenced by factors, such as genetic polymorphisms and gut microbiota composition. For instance, only certain individuals produce equol, a metabolite of daidzein with higher estrogenic potency, which may enhance or mitigate phytohormonal effects. This variability complicates the establishment of universal dietary recommendations.[31]

Implications for dietary recommendations

The dual nature of soy phytohormones – offering both benefits and risks – necessitates a cautious approach to dietary recommendations. Moderate soy consumption, such as incorporating soy milk, tofu, or soy-based snacks, is generally considered safe for adolescents. Setchell and Clerici (2010) suggest that dietary soy intake should remain within 30–50 mg of isoflavones per day to balance health benefits with minimal risk.

Parents and healthcare professionals should be aware of potential interactions between soy phytohormones and hormonal therapies or endocrine-disrupting compounds. For adolescents with existing hormonal disorders, consultation with a healthcare provider is essential before introducing soy-rich diets.

Future research should focus on long-term studies to better understand the impact of soy phytohormones on pubertal development and overall health. In addition, public health strategies should promote awareness of phytohormonal effects, enabling informed dietary choices.

Soy phytohormones present a unique opportunity for supporting hormonal regulation and development during adolescence. While their benefits in improving menstrual health and bone density are evident, potential risks associated with overconsumption warrant careful dietary planning. By fostering a balanced approach, soy phytohormones can play a valuable role in adolescent nutrition, paving the way for healthy growth and development.[32]

Table 1: Summary of key findings on soy phytohormones and adolescent development

Aspect	Key findings
Phytohormonal composition	Soybeans are rich in isoflavones, including genistein and daidzein, which structurally mimic human estrogens.
Mechanism of action	Isoflavones interact with estrogen receptors alpha and beta may exert weak estrogenic or anti-estrogenic effects depending on the hormonal milieu.
Impact on early puberty	Regular soy consumption has been associated with delayed onset of puberty in some pre-clinical studies.
Gender-specific effects	Male adolescents exhibit minimal estrogenic effects from soy, whereas females show more pronounced changes in hormonal balance.
Potential health benefits	May reduce the risk of hormonal disorders, improve bone density, and support cardiovascular health.
Risks of overconsumption	High soy intake may disrupt endocrine function or interfere with thyroid hormone synthesis in susceptible individuals.
Present research limitations	Lack of large-scale, long-term studies to confirm safety and efficacy in adolescents.

EMERGING TRENDS IN SOY-BASED HORMONAL RESEARCH

Soy-based hormonal research has gained significant traction in recent years, driven by advances in biotechnology, personalized nutrition, and therapeutic applications. The bioactive compounds in soy, particularly isoflavones, such as genistein and daidzein, exhibit potential for modulating endocrine functions and addressing various health concerns. This section explores the latest developments in soy-based hormonal research, focusing on advancements in soy enrichment, personalized nutrition, and therapeutic interventions.[33]

Advances in biotechnology for soy enrichment

Biotechnology has revolutionized the agricultural and pharmaceutical industries, and soy enrichment is no exception. Modern biotechnological approaches aim to enhance the phytohormonal content in soybeans, increasing their efficacy as dietary supplements and functional foods. Genetic engineering techniques, such as CRISPR-Cas9 and transgenic modifications, have been employed to selectively upregulate the biosynthesis of isoflavones, ensuring higher concentrations of genistein and daidzein in soybean cultivars.

Fermentation techniques have also emerged as a promising method for enriching soy-based products. Across microbial fermentation, soybeans are transformed into products with increased bioavailability of phytohormones. Fermented soy products, such as natto and miso, are particularly rich in equol, a potent metabolite of daidzein known for its estrogenic properties. These advancements make it possible to develop targeted dietary solutions for individuals with specific hormonal needs.

In addition, advancements in nanotechnology have enabled the encapsulation of soy phytohormones for enhanced delivery and stability. Nanoformulations protect isoflavones from degradation during digestion, improving their absorption and efficacy in the human body. These innovations have broadened the scope of soy-based hormonal research, paving the way for novel applications.[34]

Potential applications in personalized nutrition

The concept of personalized nutrition has gained momentum, emphasizing tailored dietary strategies based on individual genetic, metabolic, and lifestyle factors. Soy phytohormones, with their selective ER modulatory effects, hold significant potential in this domain. Research indicates that individual responses to soy-based interventions vary, influenced by genetic polymorphisms and gut microbiota composition.

One emerging area of interest is the development of nutrigenomic profiles that predict an individual's response to soy phytohormones. These profiles can guide the formulation of personalized diets, ensuring optimal hormonal modulation without adverse effects. For example, individuals who produce equol may benefit more from soy-based diets, as equol exhibits higher bioactivity than its precursor daidzein.

Another promising application is in managing hormonal imbalances and conditions, such as polycystic ovary syndrome and estrogen dominance. By leveraging phytohormones' ability to mimic or modulate natural hormones, personalized nutrition plans can address specific endocrine disorders effectively. As technology advances, soy-based nutraceuticals are likely to become an integral part of individualized dietary regimens.[35]

Use of phytohormones in therapeutic interventions

Beyond nutrition, soy phytohormones have demonstrated potential in therapeutic contexts, particularly in addressing hormonal disorders and age-related conditions. Isoflavones'

structural similarity to estrogen allows them to act as natural alternatives to hormone replacement therapy for managing menopausal symptoms. Clinical studies have shown that soy isoflavones alleviate hot flashes, improve bone density, and reduce the risk of cardiovascular diseases in postmenopausal women.

Emerging research also highlights the potential of phytohormones in cancer prevention and treatment. Genistein, in particular, has been studied for its anti-cancer properties, including its ability to inhibit tumor growth and angiogenesis in hormone-dependent cancers, such as breast and prostate cancer, by modulating estrogen and androgen pathways; soy phytohormones offer a natural and less invasive therapeutic option for such conditions.

Furthermore, soy-derived phytohormones are being explored as adjuncts in managing endocrine-related developmental disorders in adolescents. For example, their ability to balance estrogen and androgen levels can help mitigate early or delayed onset of puberty. However, more extensive clinical trials are required to establish their safety and efficacy in this population.

The field of soy-based hormonal research is rapidly evolving, driven by advances in biotechnology, personalized nutrition, and therapeutic applications. Enhanced enrichment techniques, such as genetic engineering and fermentation, have made soy phytohormones more accessible and effective. Personalized nutrition strategies leveraging individual responses to phytohormones can revolutionize dietary interventions for hormonal balance. In addition, the therapeutic potential of soy phytohormones in managing menopause, cancer, and developmental disorders underscores their versatility. As research progresses, these trends are expected to redefine the role of soy phytohormones in health and wellness.[36]

CHALLENGES AND FUTURE DIRECTIONS

Research on soy-derived phytohormones has advanced significantly, offering promising insights into their role in modulating human health, particularly during adolescence. However, the field is not without its challenges. Addressing present knowledge gaps, conducting long-term safety studies, and exploring public health integration are critical for harnessing the full potential of soy phytohormones. This section discusses these challenges and outlines future research directions.

Addressing gaps in present knowledge

Despite growing interest, there are notable gaps in the understanding of soy phytohormones' mechanisms of action and their effects on human development. While studies highlight the influence of isoflavones, such as genistein and daidzein, on endocrine pathways, inconsistencies remain in findings regarding their systemic effects. For instance, the variability in individual responses to soy phytohormones, influenced by genetic predisposition, gut microbiota composition, and environmental exposures, requires deeper investigation.

Another challenge is the lack of robust data on how phytohormones interact with other dietary components. Synergistic or antagonistic interactions between soy-derived compounds and other nutrients may significantly impact their bioactivity. Furthermore, there is limited research on gender-specific effects, especially during critical developmental stages, such as adolescence. Addressing these knowledge gaps will require interdisciplinary studies that incorporate endocrinology, nutrition, and genetics.

Long-term studies on safety and efficacy

Safety and efficacy concerns regarding soy phytohormones remain a contentious topic, particularly their long-term impact on adolescents. While short-term studies suggest benefits in hormonal regulation and the development of secondary sexual characteristics, the potential risks associated with prolonged soy consumption are not fully understood.

One key concern is the risk of hormonal imbalances due to overconsumption, potentially leading to conditions, such as estrogen dominance or disrupted androgen activity. In addition, some studies have raised concerns about the potential for soy isoflavones to mimic estrogen and contribute to hormone-dependent conditions, such as breast cancer, though evidence remains inconclusive.

To address these concerns, long-term, large-scale clinical trials are necessary. Such studies should evaluate the chronic effects of soy phytohormones on endocrine health, growth patterns, and overall well-being in adolescents. Standardizing methodologies across studies will also be critical to ensure comparability and reliability of findings.[37]

Prospects for integrating soy phytohormones in public health

The integration of soy phytohormones into public health strategies represents an exciting frontier. With the global rise in hormonal disorders and nutrition-related health challenges, soy-based interventions could serve as accessible and cost-effective solutions. For example, fortified soy products could be introduced into school meal programs to support adolescent development, particularly in regions where malnutrition and hormonal imbalances are prevalent.

However, public health initiatives must balance the benefits of soy phytohormones with potential risks. Clear dietary guidelines, tailored to different age groups and genders,

are essential to ensure safe and effective consumption. Educational campaigns highlighting the role of soy phytohormones in nutrition and health can promote informed decision-making among consumers.

The role of policymakers is also pivotal in fostering research and regulation. Governments can fund studies exploring the public health implications of soy phytohormones and establish frameworks to regulate the production and marketing of soy-based products. Collaborations between researchers, industry stakeholders, and public health officials will be essential for translating scientific findings into practical applications.

While soy phytohormones hold immense potential for improving adolescent health and addressing hormonal imbalances, significant challenges remain. Bridging knowledge gaps, conducting long-term safety studies, and developing evidence-based public health strategies are critical steps forward. By addressing these challenges, researchers can unlock new opportunities for integrating soy phytohormones into personalized nutrition and therapeutic interventions, ultimately contributing to better health outcomes on a global scale.[38]

CONCLUSION

The exploration of soybean-derived phytohormones and their influence on secondary sexual characteristics during adolescence reveals a nuanced intersection of nutrition, endocrinology, and public health. This review has synthesized findings from biochemical, clinical, and pre-clinical studies to offer a comprehensive understanding of the potential benefits and risks associated with soy phytohormones during a critical developmental period.

Key insights from the review

The key takeaway from this review is the demonstrated ability of soy-derived phytohormones, such as isoflavones (genistein and daidzein), to interact with human endocrine systems. These compounds can mimic or modulate estrogen and androgen activity, affecting secondary sexual characteristics in adolescents. The review highlights the potential of soy isoflavones to support hormonal balance, particularly in cases of hormonal imbalances or nutritional deficiencies.

In addition, comparative analyses suggest that soy-derived phytohormones offer a natural and less invasive alternative to synthetic hormonal therapies. However, variability in individual responses due to genetic, environmental, and dietary factors complicates the establishment of universal guidelines for soy consumption.

While pre-clinical studies in animals provide robust evidence of hormonal modulation, translating these findings to humans requires caution. Human studies, though promising, reveal inconsistencies stemming from variations in study design, sample size, and dietary conditions. This underscores the need for standardized methodologies and long-term studies to validate these findings further.

Implications for early adolescence and pubertal health

Adolescence represents a critical window of development, during which hormonal changes drive physical, emotional, and cognitive maturation. The role of soy phytohormones in modulating these processes has significant implications for adolescent health.

One promising application lies in addressing early-onset puberty, a condition increasingly prevalent due to lifestyle and environmental factors. Soy phytohormones could act as modulatory agents, slowing or normalizing pubertal progression by interacting with endocrine pathways. Similarly, for adolescents experiencing delayed puberty, these compounds may provide a supportive role by gently stimulating hormonal activity.

Beyond hormonal regulation, the nutritional profile of soy contributes to overall health, offering benefits, such as improved bone density, cardiovascular health, and immune function. This positions soy as a multifaceted dietary intervention with both hormonal and systemic benefits.

However, concerns regarding overconsumption and potential hormonal imbalances necessitate cautious application. Public health strategies must include education on appropriate soy intake levels and highlight the importance of a balanced diet to avoid over-reliance on phytohormones.

Final thoughts on future applications

The future of soy phytohormone research and application is promising but requires careful navigation. Advances in biotechnology, such as the development of soy products enriched with specific phytohormones, open new avenues for personalized nutrition and therapeutic interventions. For instance, tailored soy-based formulations could address specific hormonal needs in adolescents, offering targeted support during pubertal development.

Moreover, the integration of soy phytohormones into public health programs could address nutritional deficiencies in resource-limited settings. Fortified soy products in school meals or community health initiatives could help bridge gaps in adolescent nutrition, particularly in regions where access to diverse dietary options is limited.

Research gaps identified in this review, including the need for gender-specific studies and long-term clinical trials, present

opportunities for advancing the field. Interdisciplinary collaborations among nutritionists, endocrinologists, and public health experts will be critical for developing safe and effective applications of soy phytohormones.

In conclusion, soy-derived phytohormones hold significant potential for improving adolescent health by supporting hormonal regulation and addressing nutritional challenges. While promising, these applications must be guided by robust scientific evidence to ensure safety, efficacy, and equitable access. By addressing existing challenges and pursuing targeted research, the integration of soy phytohormones into dietary and therapeutic frameworks could contribute to healthier developmental outcomes for adolescents worldwide.

REFERENCES

1. Antinozzi C, Greco EA, Sgrò P, Dimauro I, Aversa A, DI Luigi L. Effects of tadalafil on skeletal muscle tissue: Exploring interactions and novel mechanisms of action. Minerva Endocrinol (Torino) 2023;48:222-9.
2. Patel J, Kumar GS, Roy H, Maddiboyina B, Leporatti S, Bohara RA. From nature to nanomedicine: Bioengineered metallic nanoparticles bridge the gap for medical applications. Discov Nano 2024;19:85.
3. Patel J, Reddy V, Kumar GS, Satyasai D, Bajari B. Gas chromatography and mass spectroscopy analysis of bioactive components on the leaf extract of *Terminalia coriacea*: A potential folklore medicinal plant. Int J Green Pharm 2017;11:S140-3.
4. Patel J, Reddy V, Kumar GS. Preliminary phytochemical screening and hepatoprotective activity of methanol extract of *Artocarpus hirsutus* leaves. Int J Phytomed 2016;8:379.
5. Zhu Y, Chen G, Diao J, Wang C. Recent advances in exploring and exploiting soybean functional peptides-a review. Front Nutr 2023;10:1185047.
6. Shaffique S, Kang SM, Hoque MIU, Imran M, Aaqil Khan M, Lee IJ. Research progress in soybean by phytohormone modulation and metal chelation over the past decade. Agriculture 2023;13:1325.
7. Maruyama K, Urano K, Kusano M, Sakurai T, Takasaki H, Kishimoto M, *et al*. Metabolite/phytohormone-gene regulatory networks in soybean organs under dehydration conditions revealed by integration analysis. Plant J 2020;103:197-211.
8. Lay AA, Fujimori E, Duarte LS, Borges AL. Prevalence and correlates of early sexual initiation among Brazilian adolescents. PLoS One 2021;16:e0260815.
9. Tuchkina IO, Pylypenko NS. Reproductive potential assesment of adolescent girls with gynecological pathology, born low to gestational weight. Mod Pediatr Ukr 2022;1:42-8.
10. Mesele J, Alemayehu A, Demissee A, Yusuf M, Abubeker F, Ahmed M, et al. Level and determinants of knowledge, attitude, and practice of risky sexual behavior among adolescents in Harar, Ethiopia. SAGE Open Med 2023;11. https://doi.org/10.1177/20503121221145539
11. Ajiboye TT, Ajiboye TO, Babalola OO. Impacts of binary oxide nanoparticles on the soybean plant and its rhizosphere, associated phytohormones, and enzymes. Molecules 2023;28:1326.
12. Llanes A, Iparraguirre J, Masciarelli O, Maria N, Luna V. Foliar application of phytohormones enhances growth of maize and soybean seedlings. Rev Invest Agropecuarias 2019;45:61-6.
13. Meng Y, Chen F, Shuai H, Luo X, Ding J, Tang S, *et al*. Karrikins delay soybean seed germination by mediating abscisic acid and gibberellin biogenesis under shaded conditions. Sci Rep 2016;6:22073.
14. Xiao CW, Wood C, Cunningham LA, Lalande M, Riding M. Effects of dietary active soybean trypsin inhibitors on pancreatic weights, histology and expression of STAT3 and receptors for androgen and estrogen in different tissues of rats. Mol Biol Rep 2021;48:4591-600.
15. Lund TD, Munson DJ, Haldy ME, Setchell KD, Lephart ED, Handa RJ. Equol is a novel anti-androgen that inhibits prostate growth and hormone feedback. Biol Reprod 2004;70:1188-95.
16. Cline JM, Söderqvist G, Register TC, Williams JK, Adams MR, Von Schoultz B. Assessment of hormonally active agents in the reproductive tract of female nonhuman primates. Toxicol Pathol 2001;29:84-90.
17. Calcaterra V, Regalbuto C, Mercurio G, Tenuta E, Verduci E, Zuccotti G, *et al*. Phytoestrogens in soybean and precocious pubertal development: Causality or casuality? Int J Pediatr Child Health. 2021;9:40-4.
18. Phiri M, Musonda E, Shasha L, Kanyamuna V, Lemba M. Individual and Community-level factors associated with early marriage in Zambia: A mixed effect analysis. BMC Womens Health 2023;23:21.
19. Palermo T, Chzhen Y, Balvin N, Kajula L, Tanzania Adolescent Cash Plus Evaluation Team. Examining determinants of gender attitudes: Evidence among Tanzanian adolescents. BMC Womens Health 2020;20:195.
20. Ojiho IH, Isokon B, Tiku TO. Family characteristics and adolescent sexual behaviour in Yala local government area of cross river state, Nigeria. Asian Res J Arts Soc Sci 2022;16:30-8.
21. Thangavel P, Puga-Olguín A, Rodríguez-Landa JF, Zepeda RC. Genistein as potential therapeutic candidate for menopausal symptoms and other related diseases. Molecules 2019;24:3892.
22. Domínguez-López I, Yago-Aragón M, Salas-Huetos A, Tresserra-Rimbau A, Hurtado-Barroso S. Effects of dietary phytoestrogens on hormones throughout a human lifespan: A review. Nutrients 2020;12:2456.
23. Suen AA, Kenan AC, Williams CJ. Developmental exposure to phytoestrogens found in soy: New findings and clinical implications. Biochem Pharmacol 2022;195:114848.
24. Mun BG, Kim HH, Yuk HJ, Hussain A, Loake GJ,

Yun BW. A potential role of coumestrol in soybean leaf senescence and its interaction with phytohormones. Front Plant Sci 2021;12:756308.
25. Tavakolan M, Alkharouf NW. Data Mining of Genomic Data Generated from Soybean Treated with Different Phytohormones. In: 2016 IEEE/ACIS 14th International Conference on Software Engineering Research, Management and Applications, SERA 2016; 2016.
26. Studham ME, MacIntosh GC. Phytohormone signaling pathway analysis method for comparing hormone responses in plant-pest interactions. BMC Res Notes 2012;5:392.
27. Medeiros RC, Santos IK, Damasceno De Goes CJ, Vasconcelos De Oliveira AL, Medeiros JA, Cobucci RN, *et al*. A comparative analysis of physical fitness of children and adolescents with HIV infection: A systematic review protocol. Medicine (Baltimore) 2019;98:e18206.
28. Wadhave AA, Jadhav AI, Arsul VA. Plant proteins applications: A review. World J Pharm Pharm Sci 2014;3:702-12.
29. Wilk M. Soja źródłem cennych składników żywieniowych. Zywnosc Nauka Technol Jakosc/Food Sci Technol Qual 2017;111:16-25.
30. Emmanuelle NE, Marie-Cécile V, Florence T, Jean-François A, Françoise L, Coralie F, *et al*. Critical role of estrogens on bone homeostasis in both male and female: From physiology to medical implications. Int J Mol Sci 2021;22:1568.
31. Satpathi S, Gaurkar SS, Potdukhe A, Wanjari MB. Unveiling the role of hormonal imbalance in breast cancer development: A comprehensive review. Cureus 2023;15:e41737.
32. Wessinger C, Hafer-Macko C, S Ryan A. Vitamin K intake in chronic stroke: Implications for dietary recommendations. Nutrients 2020;12:3059.
33. Dinsdale EC, Ward WE. Early exposure to soy isoflavones and effects on reproductive health: A review of human and animal studies. Nutrients 2010;2:1156-87.
34. Onishi H, Fuchi H, Konomi K, Hidaka O, Kamekura M. Isolation and distribution of a variety of halophilic bacteria and their classification by salt-response. Agric Biol Chem 1980;44:1253-8.
35. Merino J, Linenberg I, Bermingham KM, Ganesh S, Bakker E, Delahanty LM, *et al*. Validity of continuous glucose monitoring for categorizing glycemic responses to diet: Implications for use in personalized nutrition. Am J Clin Nutr 2022;115:1569-76.
36. Haudum C, Lindheim L, Ascani A, Trummer C, Horvath A, Münzker J, *et al*. Impact of short-term isoflavone intervention in polycystic ovary syndrome (PCOS) patients on microbiota composition and metagenomics. Nutrients 2020;12:1622.
37. Kondo T, Mizuno Y, Japanese Istradefylline Study Group. A long-term study of istradefylline safety and efficacy in patients with Parkinson disease. Clin Neuropharmacol 2015;38:41-6.
38. Levasseur W, Perré P, Pozzobon V. A review of high value-added molecules production by microalgae in light of the classification. Biotechnol Adv 2020;41:107545.

Source of Support: Nil. **Conflicts of Interest:** None declared.